

~~10484927~~

10/740 206

Connecting via Winsock to STN

Welcome to STN International! Enter x:X

LOGINID:sssptal623hrr

PASSWORD:

TERMINAL (ENTER 1, 2, 3, OR ?):2

\* \* \* \* \* Welcome to STN International \* \* \* \* \*

NEWS	1		Web Page URLs for STN Seminar Schedule - N. America
NEWS	2		"Ask CAS" for self-help around the clock
NEWS	3	Jul 12	BEILSTEIN enhanced with new display and select options, resulting in a closer connection to BABS
NEWS	4	Jul 30	BEILSTEIN on STN workshop to be held August 24 in conjunction with the 228th ACS National Meeting
NEWS	5	AUG 02	IFIPAT/IFIUDB/IFICDB reloaded with new search and display fields
NEWS	6	AUG 02	CAplus and CA patent records enhanced with European and Japan Patent Office Classifications
NEWS	7	AUG 02	The Analysis Edition of STN Express with Discover! (Version 7.01 for Windows) now available
NEWS	8	AUG 04	Pricing for the Save Answers for SciFinder Wizard within STN Express with Discover! will change September 1, 2004
NEWS	9	AUG 27	BIOCOMMERCE: Changes and enhancements to content coverage
NEWS	10	AUG 27	BIOTECHABS/BIOTECHDS: Two new display fields added for legal status data from INPADOC
NEWS	11	SEP 01	INPADOC: New family current-awareness alert (SDI) available
NEWS	12	SEP 01	New pricing for the Save Answers for SciFinder Wizard within STN Express with Discover!
NEWS	13	SEP 01	New display format, HITSTR, available in WPIDS/WPINDEX/WPIX
NEWS	14	SEP 14	STN Patent Forum to be held October 13, 2004, in Iselin, NJ
NEWS	15	SEP 27	STANDARDS will no longer be available on STN
NEWS	16	SEP 27	SWETSCAN will no longer be available on STN
NEWS EXPRESS	JULY 30 CURRENT WINDOWS VERSION IS V7.01, CURRENT MACINTOSH VERSION IS V6.0c(ENG) AND V6.0Jc(JP), AND CURRENT DISCOVER FILE IS DATED 11 AUGUST 2004		
NEWS HOURS	STN Operating Hours Plus Help Desk Availability		
NEWS INTER	General Internet Information		
NEWS LOGIN	Welcome Banner and News Items		
NEWS PHONE	Direct Dial and Telecommunication Network Access to STN		
NEWS WWW	CAS World Wide Web Site (general information)		

Enter NEWS followed by the item number or name to see news on that specific topic.

All use of STN is subject to the provisions of the STN Customer agreement. Please note that this agreement limits use to scientific research. Use for software development or design or implementation of commercial gateways or other similar uses is prohibited and may result in loss of user privileges and other penalties.

\* \* \* \* \* STN Columbus \* \* \* \* \*

FILE 'HOME' ENTERED AT 14:36:53 ON 28 SEP 2004

10484927

=>

Uploading

THIS COMMAND NOT AVAILABLE IN THE CURRENT FILE

Do you want to switch to the Registry File?

Choice (Y/n):

Switching to the Registry File...

Some commands only work in certain files. For example, the EXPAND command can only be used to look at the index in a file which has an index. Enter "HELP COMMANDS" at an arrow prompt (=>) for a list of commands which can be used in this file.

=> FILE REGISTRY

COST IN U.S. DOLLARS	SINCE FILE ENTRY	TOTAL SESSION
FULL ESTIMATED COST	0.21	0.21

FILE 'REGISTRY' ENTERED AT 14:37:03 ON 28 SEP 2004

USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.

PLEASE SEE "HELP USAGETERMS" FOR DETAILS.

COPYRIGHT (C) 2004 American Chemical Society (ACS)

Property values tagged with IC are from the ZIC/VINITI data file provided by InfoChem.

STRUCTURE FILE UPDATES: 27 SEP 2004 HIGHEST RN 752974-11-1

DICTIONARY FILE UPDATES: 27 SEP 2004 HIGHEST RN 752974-11-1

TSCA INFORMATION NOW CURRENT THROUGH MAY 21, 2004

Please note that search-term pricing does apply when conducting SmartSELECT searches.

Crossover limits have been increased. See HELP CROSSOVER for details.

Experimental and calculated property data are now available. For more information enter HELP PROP at an arrow prompt in the file or refer to the file summary sheet on the web at:  
<http://www.cas.org/ONLINE/DBSS/registryss.html>

=>


Uploading C:\STNEXP4\QUERIES\10740206.str

L1 STRUCTURE UPLOADED

=> d l1

L1 HAS NO ANSWERS

L1 STR



10484927

Structure attributes must be viewed using STN Express query preparation.

=> s l1 full

FULL SEARCH INITIATED 14:37:22 FILE 'REGISTRY'  
FULL SCREEN SEARCH COMPLETED - 68940 TO ITERATE

100.0% PROCESSED 68940 ITERATIONS  
SEARCH TIME: 00.00.02

183 ANSWERS

L2 183 SEA SSS FUL L1

=> file ca

COST IN U.S. DOLLARS

SINCE FILE

TOTAL

ENTRY

SESSION

FULL ESTIMATED COST

155.84

156.05

FILE 'CA' ENTERED AT 14:38:00 ON 28 SEP 2004

USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.

PLEASE SEE "HELP USAGETERMS" FOR DETAILS.

COPYRIGHT (C) 2004 AMERICAN CHEMICAL SOCIETY (ACS)

Copyright of the articles to which records in this database refer is held by the publishers listed in the PUBLISHER (PB) field (available for records published or updated in Chemical Abstracts after December 26, 1996), unless otherwise indicated in the original publications. The CA Lexicon is the copyrighted intellectual property of the American Chemical Society and is provided to assist you in searching databases on STN. Any dissemination, distribution, copying, or storing of this information, without the prior written consent of CAS, is strictly prohibited.

FILE COVERS 1907 - 23 Sep 2004 VOL 141 ISS 14

FILE LAST UPDATED: 23 Sep 2004 (20040923/ED)

This file contains CAS Registry Numbers for easy and accurate substance identification.

=> s l2

L3 92 L2

=> s l3 and salt

714335 SALT

L4 9 L3 AND SALT

=> s l3 and sulfonate

52599 SULFONATE

L5 0 L3 AND SULFONATE

=> d l4 1-9 sub bib abs

'SUB' IS NOT A VALID FORMAT FOR FILE 'CA'

The following are valid formats:

ABS ----- GI and AB

ALL ----- BIB, AB, IND, RE

APPS ----- AI, PRAI

BIB ----- AN, plus Bibliographic Data and PI table (default)

CAN ----- List of CA abstract numbers without answer numbers

CBIB ----- AN, plus Compressed Bibliographic Data


10484927

DALL ----- ALL, delimited (end of each field identified)  
DMAX ----- MAX, delimited for post-processing  
FAM ----- AN, PI and PRAI in table, plus Patent Family data  
FBIB ----- AN, BIB, plus Patent FAM  
IND ----- Indexing data  
IPC ----- International Patent Classifications  
MAX ----- ALL, plus Patent FAM, RE  
PATS ----- PI, SO  
SAM ----- CC, SX, TI, ST, IT  
SCAN ----- CC, SX, TI, ST, IT (random display, no answer numbers;  
SCAN must be entered on the same line as the DISPLAY,  
e.g., D SCAN or DISPLAY SCAN)  
STD ----- BIB, IPC, and NCL  
  
IABS ----- ABS, indented with text labels  
IALL ----- ALL, indented with text labels  
IBIB ----- BIB, indented with text labels  
IMAX ----- MAX, indented with text labels  
ISTD ----- STD, indented with text labels  
  
OBIB ----- AN, plus Bibliographic Data (original)  
OIBIB ----- OBIB, indented with text labels  
  
SBIB ----- BIB, no citations  
SIBIB ----- IBIB, no citations  
  
HIT ----- Fields containing hit terms  
HITIND ----- IC, ICA, ICI, NCL, CC and index field (ST and IT)  
containing hit terms  
HITRN ----- HIT RN and its text modification  
HITSTR ----- HIT RN, its text modification, its CA index name, and  
its structure diagram  
HITSEQ ----- HIT RN, its text modification, its CA index name, its  
structure diagram, plus NTE and SEQ fields  
FHITSTR ----- First HIT RN, its text modification, its CA index name, and  
its structure diagram  
FHITSEQ ----- First HIT RN, its text modification, its CA index name, its  
structure diagram, plus NTE and SEQ fields  
KWIC ----- Hit term plus 20 words on either side  
OCC ----- Number of occurrence of hit term and field in which it occurs

To display a particular field or fields, enter the display field codes. For a list of the display field codes, enter HELP DFIELDS at an arrow prompt (=>). Examples of formats include: TI; TI,AU; BIB,ST; TI,IND; TI,SO. You may specify the format fields in any order and the information will be displayed in the same order as the format specification.

All of the formats (except for SAM, SCAN, HIT, HITIND, HITRN, HITSTR, FHITSTR, HITSEQ, FHITSEQ, KWIC, and OCC) may be used with DISPLAY ACC to view a specified Accession Number.  
ENTER DISPLAY FORMAT (BIB):abs bib

L4 ANSWER 1 OF 9 CA COPYRIGHT 2004 ACS on STN  
AB The invention relates to a method for the identification of enzymes that are preferentially expressed in certain tumor tissue as compared with rapidly growing normal cells or tissue and the use of the enzymes to design compds. which generate active anticancer substances selectively in tumor tissue. Compds. X-Y-Q [X is a pro-moiety that is designed to generate an active anticancer substance (Q-Y-H) selectively in tumors by the enzymes; Q-Y- is a radical derived from the active anticancer substance in which Y is O, S or N] and their pharmaceutically-acceptable



10484927



salts are claimed. Thus, 13 $\alpha$ -[(2R,3S)-2-[(5S)-[5-[(2S)-(2-aminopropionyl)amino]-5-hydroxycarbonyl]pentanoyloxy]-3-(benzoylamino)-3-phenylpropionyloxy]-2a-(benzyloxy)-4a,10 $\beta$ -diacetoxy-1 $\beta$ ,7 $\beta$ -dihydroxy-5 $\beta$ ,20-epoxytax-1-en-9-one formic acid salt (I) was prepared by reaction of taxol with (2S)-2-[(2S)-2-(benzyloxycarbonylamino)-3-phenylpropionylamino]hexanedioic acid 1-benzyl ester. Compound I showed cytotoxicity IC<sub>50</sub> = 51 nM after 24 h against human colon cancer cell line HCT116.

AN 139:7174 CA  
TI Method for identification of tumor targeting enzymes for design of compounds which generate anticancer substances  
IN Ishitsuka, Hideo; Okabe, Hisafumi; Shimma, Nobuo; Tsukuda, Takuo; Umeda, Isao  
PA F. Hoffmann-La Roche A.-G., Switz.  
SO PCT Int. Appl., 118 pp.  
CODEN: PIXXD2  
DT Patent  
LA English  
FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	WO 2003043631	A2	20030530	WO 2002-EP12911	20021118
	W:	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM			
	RW:	GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG			
	US 2003138864	A1	20030724	US 2002-301460	20021121
PRAI	EP 2001-127401	A	20011123		
	EP 2001-130245	A	20011219		
	EP 2002-5298	A	20020312		
OS	MARPAT 139:7174				

L4 ANSWER 2 OF 9 CA COPYRIGHT 2004 ACS on STN  
AB Motuporin (I) was prepared by a convergent synthesis in which all stereocenters are derived from common amino acids or from D-mandelic acid; the 3 unusual amino acids in I are all derived from D-threonine.  
AN 124:30333 CA  
TI Enantiospecific total synthesis of the protein phosphatase inhibitor motuporin.  
AU Valentekovich, Robert J.; Schreiber, Stuart L.  
CS Department of Chemistry, Harvard University, Cambridge, MA, 02138, USA  
SO Journal of the American Chemical Society (1995), 117(35), 9069-70  
CODEN: JACSAT; ISSN: 0002-7863  
PB American Chemical Society  
DT Journal  
LA English  
OS CASREACT 124:30333

L4 ANSWER 3 OF 9 CA COPYRIGHT 2004 ACS on STN  
GI



Ser(Bzl)-Asn-Leu-Ser(Bzl)-Thr(Bzl)-Asu-Val-Leu-  
 Gly-Lys(Cl-Z)-Leu-Ser(Bzl)-Gln-Glu(OBzl)-Leu-  
 His(Tos)-Lys(Cl-Z)-Leu-Gln-Thr(Bzl)-Tyr(Br-Z)-  
 Pro-Arg(Tos)-Thr(Bzl)-Asp(OBzl)-Val-Gly-Ala-Gly-  
 Thr(Bzl)-Pro-Resin

IV

AB Cyclopeptides (I; X1 = OH, NH<sub>2</sub>, amino acid or peptide residue, wherein each amino acid residue may be protected) are prepared by cyclization of resin-bound open chain peptides [II; A = Ser(X1)-Asn-Leu-Ser(X1)-Thr(X1)-X<sub>2</sub>, Ser(X1)-Asn-Leu-Ser(X1)-X<sub>2</sub>, Ser(X1)-Asn-Leu-X<sub>2</sub>, Ser(X1)-Asn-X<sub>2</sub>, Ser(X1)-X<sub>2</sub>; B = X<sub>3</sub>, X<sub>3</sub>-Thr(X1), X<sub>3</sub>-Ser(X1)-Thr(X1), X<sub>3</sub>-Leu-Ser(X1)-Thr(X1), X<sub>3</sub>-Asn-Leu-Ser(X1)-Thr(X1); X<sub>1</sub> = H, HO-protective group; X<sub>3</sub> = H, H<sub>2</sub>N-protective group; X = direct bond to Resin, amino acid or peptide residue; Resin = resin for solid phase reaction] without using protease followed by resin cleavage. The cyclization proceeds in high yields without side reactions and gives calcitonin derivs., particularly elcatonin, in good yields. Thus, 620 mg R-Leu-Ser(Bzl)-Thr(Bzl)-Asu[Ser(Bzl)-Asn-OR<sub>1</sub>]-Val-Leu-Gly-Lys(Cl-Z)-Leu-Ser(Bzl)-Gln-Glu(OBzl)-Leu-His(Tos)-Lys(Cl-Z)-Leu-Gln-Thr(Bzl)-Tyr(Br-Z)-Pro-Arg(Tos)-Thr(Bzl)-Asp(OBzl)-Val-Gly-Ala-Gly-Thr(Bzl)-Pro-p-methylbenzhydrylamine polystyrene resin (III; R = Boc, Asu = 2-aminosuberic acid residue, R<sub>1</sub> = CMe<sub>3</sub>, Bzl = CH<sub>2</sub>Ph, Cl-Z = 2-chlorobenzoyloxycarbonyl) (preparation given) was stirred with 50% CF<sub>3</sub>CO<sub>2</sub>H in CH<sub>2</sub>Cl<sub>2</sub> at room temperature for 30 min followed by washing with CH<sub>2</sub>Cl<sub>2</sub>, 10% (Me<sub>2</sub>CH)<sub>2</sub>NEt in DMF, and DMF, III (R = R<sub>1</sub> = H) which was added to N-methylpyrrolidone and treated with 1-hydroxybenzotriazole monohydrate and DCC at room temperature for 24 h to give resin bound protected cyclopeptide (IV; Resin = p-methylbenzhydrylamine polystyrene resin). The latter peptide was treated with HF(1) and anisole at 0° for 1 h, distilled in vacuo, washed with Et<sub>2</sub>O, and extracted with 1 M aqueous AcOH solution; the extract was

lyophilized to give 260 mg crude elcatonin which was purified by ion exchange chromatog. and reversed phase HPLC using TSK Gel ODS-120 column (Toso Inc., Ltd.) and salt exchange to give elcatonin AcOH salt.

AN 122:106541 CA

TI Preparation of cyclic peptides containing  $\alpha$ -aminosuberic acid as calcitonin derivatives

IN Inoe, Takashi; Kimura, Hitoshi

PA Daicel Chem, Japan

SO Jpn. Kokai Tokkyo Koho, 15 pp.

CODEN: JKXXAF

DT Patent

LA Japanese

FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	JP 06157593	A2	19940603	JP 1992-339821	19921125
PRAI	JP 1992-339821		19921125		
OS	CASREACT 122:106541; MARPAT 122:106541				

L4 ANSWER 4 OF 9 CA COPYRIGHT 2004 ACS on STN

AB Two approaches to the synthesis of 2'-O-(N-alkylsuccinamoyl) erythromycin derivs. are explored as a means of introducing anionic and cationic functionalities into erythromycin at the 2'-O-position. The phenacyl group was used to protect the anionic functionality during the synthesis of 2'-O-[N-(carboxylatomethyl)succinamoyl] erythromycin and disodium 2'-O-[N-(1,3-dicarboxylatopropyl)succinamoyl] erythromycin, deprotection being achieved using sodium thiophenoxide under anhydrous conditions. 2'-O-(N-[2-Dimethylammonio)ethyl]succinamoyl] erythromycin lactobionate and (S)-2'-O-[N-(1-methoxycarbonyl-3-guanidinopropyl)succinamoyl] erythromycin required no protection and were prepared in an otherwise similar fashion by mixed anhydride activation of 2'-O-(3-carboxypropanoyl) erythromycin, followed by treatment with the appropriate alkylamine. Sodium 2'-O-[N-(2-sulfonatoethyl)succinamoyl] erythromycin, disodium 2'-O-[N-(1-carboxylato-2-sulfonatoethyl)succinamoyl] erythromycin and trisodium 2'-O-[N-(1-carboxylato-2-phosphonatoethyl)succinamoyl] erythromycin were prepared in a different fashion by treatment of activated 2'-O-(3-carboxypropanoyl) erythromycin with the appropriate amino acid in a Schotten-Baumann related procedure. The aqueous solubilities of derivs. in 0.1 M phosphate buffer are reported along with some preliminary stability information.

AN 115:72090 CA

TI Approaches to novel water-soluble prodrugs of erythromycin A. Synthesis of 2'-O-(N-alkylsuccinamoyl)erythromycin derivatives incorporating anionic and cationic groups

AU Ackland, Mark J.; Atkins, Paul J.; Jones, Norman B.

CS Upjohn Lab.-UK, Upjohn Ltd., Crawley/West Sussex, RH10 2NJ, UK

SO Journal of Chemical Research, Synopses (1991), (6), 142-3

CODEN: JRPSDC; ISSN: 0308-2342

DT Journal

LA English

L4 ANSWER 5 OF 9 CA COPYRIGHT 2004 ACS on STN

AB Human proinsulin C-peptide was synthesized by the solid-phase method. The product was purified consecutively by gel filtration, DEAE-cellulose chromatog., and high-performance liquid chromatog. (HPLC). The purified material behaved as a single component in reversed-phase HPLC, gave correct amino acid ratios, and was not distinguished from natural human C-peptide in terms of immunoreactivity and chromatog. behaviors. The  $\alpha \rightarrow \beta$  transpeptidation at the Asp-Leu sequence, possible to occur associated with the HF cleavage, was studied using model peptides to demonstrate that the formation of  $\beta$ -peptide was 3-4% regardless of whether the  $\beta$ -carboxylic acid is free or protected as a benzyl ester.

AN 96:52660 CA

TI A synthesis of human proinsulin C-peptide

AU Igano, Kenichi; Minotani, Yuriko; Yoshida, Nobuo; Kono, Masao; Inouye, Ken

CS Shionogi Res. Lab., Shionogi Co., Ltd., Osaka, 553, Japan

SO Bulletin of the Chemical Society of Japan (1981), 54(10), 3088-94


CODEN: BCSJA8; ISSN: 0009-2673

DT Journal

LA English

L4 ANSWER 6 OF 9 CA COPYRIGHT 2004 ACS on STN

GI



AB Peptides containing aspartic acid  $\beta$ -phenacyl ester residues, which do not cyclize under conditions of acidolysis, underwent ring closure to the corresponding succinimido derivs. under basic conditions. When the Cl<sup>+</sup> salt of BOC-Asp-Gly-NHNA (BOC = Me<sub>3</sub>CO<sub>2</sub>C, NA =  $\beta$ -naphthyl) was esterified with BrCH<sub>2</sub>COPh, succinimido peptide I (X = Gly) was obtained by cyclization of BOC-Asp(OCH<sub>2</sub>COPh)-Gly-NHNA. BOC-Asp(OCH<sub>2</sub>COPh)-Val-NHNA was also prepared and cyclized to I (X = Val) under basic conditions.

AN 89:129915 CA

TI Side reactions in peptide synthesis. 8. On the phenacyl group in the protection of the  $\beta$ -carboxyl function of aspartyl residues

AU Bodanszky, Miklos; Martinez, Jean

CS Dep. Chem., Case Western Reserve Univ., Cleveland, OH, USA

SO Journal of Organic Chemistry (1978), 43(15), 3071-3

CODEN: JOCEAH; ISSN: 0022-3263

DT Journal

LA English

L4 ANSWER 7 OF 9 CA COPYRIGHT 2004 ACS on STN

AB The 1,5-diazabicyclo[4.3.0]non-5-ene and/or 1,8-diazabicyclo[5.4.0]undec-7-ene salts of Me<sub>3</sub>CO<sub>2</sub>C-X-OH [X = Gly, Ala, Val, Ile, Tyr(CH<sub>2</sub>Ph), Arg(SO<sub>2</sub>C<sub>6</sub>H<sub>4</sub>Me-p), Gly-Ile, Trp-Gly], PhCH<sub>2</sub>O<sub>2</sub>C-Leu-Ala-OH, PhCH<sub>2</sub>O<sub>2</sub>C-Val-Phe-Gly-OH, and p-(O<sub>2</sub>N)C<sub>6</sub>H<sub>4</sub>CH<sub>2</sub>O<sub>2</sub>C-Y(OCH<sub>2</sub>C<sub>6</sub>H<sub>4</sub>NO<sub>2</sub>-p)-OH (Y = Asp, Glu) were esterified to chloromethylated styrene-divinylbenzene polymer in 59-100% yields at 50° for 28 h. The above procedure for the incorporation of N-protected peptides onto the resin did not cause racemization.

AN 87:102642 CA

TI An improved attachment of N-protected amino acid and peptide to chloromethylated polystyrene-divinylbenzene resin

AU Suzuki, Kenji; Endo, Nobuyoshi

CS Tohoku Coll. Pharm., Sendai, Japan

SO Chemical & Pharmaceutical Bulletin (1977), 25(5), 1143-6

CODEN: CPBTAL; ISSN: 0009-2363

DT Journal

LA English

L4 ANSWER 8 OF 9 CA COPYRIGHT 2004 ACS on STN

AB Different published observations of the authors (CA 56, 12153f; 58, 3645e) on the separation of carnitine (I) from possible precursors of I, including 4-aminobutyric acid (II) and some of its hydroxylated and N-methylated derivs., have been reviewed and discussed. In ion-exchange chromatography a satisfactory separation of II from I could not be obtained on a Dowex 50 resin using HCl as eluant because methylation of amino group increases the nonionic adsorption to the resin and, furthermore, the methylated amino group would be expected to be less hydrated than the unsubstituted amino group, as a consequence of which the smaller ions would be held more firmly to the resin by ionic force. Hence an ion-exchange resin with another polarity of the skeleton such as a sulfonated phenolic resin, e.g. Duolite C 3 had been recommended. Thin-layer chromatography has been found to be very convenient to use for a rapid semiquant. determination of the emerging compds. and for their identification.

AN 63:89255 CA



10484927

OREF 63:16454h,16455a-b

TI Separation of carnitine from related compounds by ion-exchange and thin-layer chromatography

AU Lindstedt, Goran; Lindstedt, Sven

CS Karolinska Inst., Stockholm

SO Recent Res. Carnitine, Its Relation Lipid Metab., Papers Symp., Cambridge, Mass. (1965), Volume Date 1964 11-21

DT Journal

LA English

L4 ANSWER 9 OF 9 CA COPYRIGHT 2004 ACS on STN

AB A mixture of 60 g. p-toluene-sulfonic acid monohydrate, 40 g. L-aspartic acid, and 100 mL. benzyl alc. was refluxed with stirring for 18 h. H<sub>2</sub>O formed was removed through a Dean-Stark separator. Cooling in ice gave 74% dibenzyl L-aspartate p-toluenesulfonate (I), m. 157-8° (aqueous EtOH),  $[\alpha]_{25D}^{20} 1^\circ$  (EtOH). Similarly prepared were dibenzyl L-glutamate p-toluenesulfonate (88%), m. 139.5-41.5° (H<sub>2</sub>O);  $\gamma$ -benzyl L-glutamate p-toluenesulfonate, m. 147-53° (EtOH-Et<sub>2</sub>O); L-leucine benzyl ester p-toluenesulfonate (62%), m. 211.5-12.5° (H<sub>2</sub>O). I (100 g.) in 500 mL. Et<sub>2</sub>O shaken with 45 g. cold Na<sub>2</sub>CO<sub>3</sub> and 100 mL. H<sub>2</sub>O gave 88% dibenzyl L-aspartate (II), oil; HCl salt (60%) m. 129.5-30° (MeOH-Et<sub>2</sub>O),  $[\alpha]_{25D}^{20} 1^\circ$  (EtOH). Likewise prepared were: dibenzyl L-glutamate HCl salt, m. 99-100°,  $[\alpha]_{25D}^{20} 9^\circ$  (HCl), and  $\gamma$ -benzyl

L-glutamate (43%), m. 162-4°,  $[\alpha]_{25D}^{20} 19^\circ$  (AcOH). II on long standing deposited 3,6-bis(benzyloxycarbonyl)-2,5-piperazinedione, m. 157-8° (EtOH). L-Aspartic acid (66.5 g.), 500 mL. benzyl alc., and 50 mL. concentrated H<sub>2</sub>SO<sub>4</sub> was kept 20 h. at room temperature and then basified

with 200 mL. C<sub>5</sub>H<sub>5</sub>N in 1 l. EtOH. The precipitate was filtered off to give 63%  $\beta$ -benzyl L-aspartate, m. 211.5-12.5° (decomposition) (aqueous C<sub>5</sub>H<sub>5</sub>N),  $[\alpha]_{25D}^{20} 27^\circ$  (HCl). Freshly prepared II (9.4 g.), 50 mL. Et<sub>2</sub>O, 1.9 g. Na<sub>2</sub>CO<sub>3</sub>, and 4.5 mL. H<sub>2</sub>O were stirred with dropwise addition of  $\beta$ -carbomethoxypropionyl chloride in 50 mL. Et<sub>2</sub>O for 30 min. at 15-17°. After addnl. stirring for 30 min., the Et<sub>2</sub>O layer was separated, dried (Na<sub>2</sub>SO<sub>4</sub>), and concentrated to give 71% N-( $\beta$ -carbomethoxypropionyl)-L-aspartic acid dibenzyl ester, m. 57-8° (petr. ether). Similarly prepared was 90% N-( $\beta$ -carbomethoxypropionyl)-L-aspartic acid dibenzyl ester (III), m. 70-1.5°. II (28.8 g.), 9 g. succinic anhydride, and 300 mL. CHCl<sub>3</sub> heated 30 min. gave 96% N-( $\beta$ -carboxypropionyl)-L-aspartic acid dibenzyl ester (IV), m. 97-8° (CCl<sub>4</sub>). III and IV on catalytic hydrogenation sep. gave N-( $\beta$ -carboxypropionyl)-L-aspartic acid, oil; solid tri-Me ester; tris(p-bromophenacyl) ester m. 138-9° (decomposition) (EtOH); triamide m. 220.5-21° (decomposition) (H<sub>2</sub>O). II (4.85 g.), 2.3 g. phthalic anhydride, 0.5 mL. Et<sub>3</sub>N, and 100 mL. PhMe were heated for 3.5 h. to give 6.3 g. dibenzyl N-phthaloyl-L-aspartate, oil, which on catalytic hydrogenation gave 3.7 g. N-phthaloyl-L-aspartic acid, m. 223-6° (H<sub>2</sub>O),  $[\alpha]_{25D}^{20} -39^\circ$  (EtOH). Use of HCONMe<sub>2</sub> or AcOH gave the DL-ester. Similarly prepared were: 68.5% N-phthaloyl-L-glutamic acid, m. 158-60° (H<sub>2</sub>O),  $[\alpha]_{25D}^{20} -45^\circ$  (EtOH); 38% N-phthaloyl-L-leucine, m. 119-21° (aqueous MeOH),  $[\alpha]_{25D}^{20} -24^\circ$  (EtOH); and 92.5% di-Et N-phthaloyl-L-aspartate,  $[\alpha]_{25D}^{20} -45^\circ$  (EtOH).

AN 60:9998 CA

OREF 60:1831d-h

TI Some derivatives of aspartic and glutamic acids

AU Bose, Ajay K.; Strube, Richard E.

CS Upjohn Co., Kalamazoo, MI

SO Journal of Pharmaceutical Sciences (1963), 52(9), 847-51

CODEN: JPMSAE; ISSN: 0022-3549

DT Journal

4328

202-2204415

10484927

LA Unavailable  
OS CASREACT 60:9998